

## Corresponding functions of SW in GSW

Here we present a table that shows some function names in the GSW Oceanographic Toolbox of TEOS-10 and the corresponding function names in the SeaWater Matlab Library of EOS-80, [http://www.cmar.csiro.au/datacentre/ext\\_docs/seawater.htm](http://www.cmar.csiro.au/datacentre/ext_docs/seawater.htm).

Variable	SeaWater & ESO-80	Gibbs-SeaWater (GSW) & TEOS-10
Absolute Salinity	-	gsw_SA_from_SP(SP,p,long,lat)
Conservative Temperature	-	gsw_CT_from_t(SA,t,p)
density (i.e. in situ density)	sw_dens(SP,t,p)	gsw_rho(SA,CT,p)
potential density	sw_pden(SP,t,p,pr)	gsw_rho(SA,CT,pr)
potential temperature	sw_ptmp(SP,t,p,pr)	gsw_pt_from_t(SA,t,p,pr)
in situ temperature from pt	sw_temp(SP,pt,p,pr)	gsw_pt_from_t(SA,pt,pr,p)
$\sigma_0$ , using $\theta_0 = \text{sw\_ptmp}(\text{SP}, \text{t}, \text{p}, 0)$	sw_dens(SP, $\theta_0$ , 0) - 1000 kg m <sup>-3</sup>	gsw_sigma0(SA,CT)
$\sigma_2$ , using $\theta_2 = \text{sw\_ptmp}(\text{SP}, \text{t}, \text{p}, 2000)$	sw_dens(SP, $\theta_2$ , 2000) - 1000 kg m <sup>-3</sup>	gsw_sigma2(SA,CT)
$\sigma_4$ , using $\theta_4 = \text{sw\_ptmp}(\text{SP}, \text{t}, \text{p}, 4000)$	sw_dens(SP, $\theta_4$ , 4000) - 1000 kg m <sup>-3</sup>	gsw_sigma4(SA,CT)
specific volume anomaly	sw_svan(SP,t,p)	gsw_specvol_anom(SA,CT,p)
dynamic height anomaly	- sw_gpan(SP,t,p)	gsw_geo_strf_dyn_height(SA,CT,p,p_ref)
geostrophic velocity	sw_gvel(ga,lat,long)	gsw_geostrophic_velocity(geo_str,long,lat,p)
$N^2$	sw_bfrq(SP, t, p, lat)	gsw_Nsquared(SA,CT,p,lat)
pressure from height (SW uses depth, not height)	sw_pres(- z,lat)	gsw_p_from_z(z,lat)
height from pressure (SW outputs depth, not height)	z = - sw_dpth(p,lat)	gsw_z_from_p(p,lat)
sound speed	sw_svel(SP,t,p)	gsw_sound_speed_CT_exact(SA,CT,p), or gsw_sound_speed(SA,CT,p), or gsw_sound_speed_t_exact(SA,t,p)
isobaric heat capacity	sw_cp(SP,t,p)	gsw_cp_t_exact(SA,t,p)
adiabatic lapse rate*	sw_adtg(SP,t,p)	gsw_adiabatic_lapse_rate_from_CT(SA,CT,p), or gsw_adiabatic_lapse_rate_from_t(SA,t,p)
SP from conductivity ratio, (PSS-78)	sw_salt(R,t,p)	gsw_SP_from_R(R,t,p)
conductivity ratio from SP, (PSS-78)	sw_cndr(SP,t,p)	gsw_R_from_SP(SP,t,p)
distance	sw_dist(lat,long,units)	gsw_distance(long,lat,p)
gravitational acceleration	sw_g(lat,z)	gsw_grav(lat,p)
Coriolis parameter	sw_f(lat)	gsw_f(lat)
testing of all functions	sw_test	gsw_check_functions
contents	Contents	gsw_contents

\* The SW and GSW functions output the adiabatic lapse rate in different units, being K (dbar)<sup>-1</sup> and K Pa<sup>-1</sup> respectively.