

Notes on the function gsw_beta_const_t_exact(S_A, t, p)

This function, **gsw_beta_const_t_exact**(S_A, t, p), evaluates the saline contraction coefficient with respect to Absolute Salinity β^θ at constant in situ temperature t , given in terms of the TEOS-10 Gibbs function by Eqn. (2.19.1) of the TEOS-10 Manual (IOC *et al.*, 2010), repeated here,

$$\beta^t = \beta^t(S_A, t, p) = \left. \frac{1}{\rho} \frac{\partial \rho}{\partial S_A} \right|_{T,p} = - \left. \frac{1}{v} \frac{\partial v}{\partial S_A} \right|_{T,p} = - \frac{g_{S_A P}(S_A, t, p)}{g_P(S_A, t, p)}. \quad (2.19.1)$$

This function uses the full TEOS-10 Gibbs function $g(S_A, t, p)$ of IOC *et al.* (2010), being the sum of the IAPWS-09 and IAPWS-08 Gibbs functions.

References

- IAPWS, 2008: Release on the IAPWS Formulation 2008 for the Thermodynamic Properties of Seawater. The International Association for the Properties of Water and Steam. Berlin, Germany, September 2008, available from www.iapws.org. This Release is referred to in the text as **IAPWS-08**.
- IAPWS, 2009: Supplementary Release on a Computationally Efficient Thermodynamic Formulation for Liquid Water for Oceanographic Use. The International Association for the Properties of Water and Steam. Doorwerth, The Netherlands, September 2009, available from <http://www.iapws.org>. This Release is referred to in the text as **IAPWS-09**.
- IOC, SCOR and IAPSO, 2010: *The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties*. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56, UNESCO (English), 196 pp. Available from <http://www.TEOS-10.org>

Here follows section 2.19 and appendix A.15 of the TEOS-10 Manual (IOC *et al.*, 2010).

2.19 Saline contraction coefficients

The saline contraction coefficient β^t (sometimes also called the haline contraction coefficient) at constant *in situ* temperature t , is

$$\beta^t = \beta^t(S_A, t, p) = \left. \frac{1}{\rho} \frac{\partial \rho}{\partial S_A} \right|_{T,p} = - \left. \frac{1}{v} \frac{\partial v}{\partial S_A} \right|_{T,p} = - \frac{g_{S_A P}}{g_P}. \quad (2.19.1)$$

The saline contraction coefficient β^θ at constant potential temperature θ , is (see appendix A.15)

$$\begin{aligned} \beta^\theta = \beta^\theta(S_A, t, p, p_r) &= \left. \frac{1}{\rho} \frac{\partial \rho}{\partial S_A} \right|_{\theta, p} = - \left. \frac{1}{v} \frac{\partial v}{\partial S_A} \right|_{\theta, p} \\ &= \frac{g_{TP} [g_{S_A T} - g_{S_A T}(S_A, \theta, p_r)] - g_{TT} g_{S_A P}}{g_P g_{TT}}, \end{aligned} \quad (2.19.2)$$

where p_r is the reference pressure of θ . One of the $g_{S_A T}$ derivatives in the numerator is evaluated at (S_A, θ, p_r) whereas all the other derivatives are evaluated at (S_A, t, p) .

The saline contraction coefficient β^Θ at constant Conservative Temperature Θ , is (see appendix A.15)

$$\begin{aligned}\beta^\Theta &= \beta^\Theta(S_A, t, p) = \frac{1}{\rho} \frac{\partial \rho}{\partial S_A} \bigg|_{\Theta, p} = - \frac{1}{v} \frac{\partial v}{\partial S_A} \bigg|_{\Theta, p} \\ &= \frac{g_{TP} \left[g_{S_A T} - (T_0 + \theta)^{-1} g_{S_A}(S_A, \theta, 0) \right] - g_{TT} g_{S_A P}}{g_P g_{TT}}.\end{aligned}\quad (2.19.3)$$

Note that Conservative Temperature Θ is defined only with respect to a reference pressure of 0 dbar as indicated in this equation. The g_{S_A} derivative in the numerator is evaluated at $(S_A, \theta, 0)$ whereas all the other derivatives are evaluated at (S_A, t, p) .

In the SIA computer software (appendix M) all three saline contraction coefficients are produced in units of kg kg^{-1} while in the GSW Oceanographic Toolbox (appendix N) all three saline contraction coefficients are produced in units of kg g^{-1} consistent with the preferred oceanographic unit for S_A being g kg^{-1} .

A.15 Derivation of the expressions for α^θ , β^θ , α^Θ and β^Θ

This appendix derives the expressions in Eqns. (2.18.2) – (2.18.3) and (2.19.2) – (2.19.3) for the thermal expansion coefficients α^θ and α^Θ and the haline contraction coefficients β^θ and β^Θ .

In order to derive Eqn. (2.18.2) for α^θ we first need an expression for $\partial\theta/\partial T|_{S_A, p}$. This is found by differentiating with respect to *in situ* temperature the entropy equality $\eta(S_A, t, p) = \eta(S_A, \theta[S_A, t, p, p_r], p_r)$ which defines potential temperature, obtaining

$$\frac{\partial\theta}{\partial T} \bigg|_{S_A, p} = \frac{\eta_T(S_A, t, p)}{\eta_T(S_A, \theta, p_r)} = \frac{g_{TT}(S_A, t, p)}{g_{TT}(S_A, \theta, p_r)} = \frac{(T_0 + \theta) c_p(S_A, t, p)}{(T_0 + t) c_p(S_A, \theta, p_r)}.\quad (A.15.1)$$

This is then used to obtain the desired expression Eqn. (2.18.2) for α^θ as follows

$$\alpha^\theta = \frac{1}{v} \frac{\partial v}{\partial \theta} \bigg|_{S_A, p} = \frac{1}{v} \frac{\partial v}{\partial T} \bigg|_{S_A, p} \left(\frac{\partial\theta}{\partial T} \bigg|_{S_A, p} \right)^{-1} = \frac{g_{TP}(S_A, t, p)}{g_P(S_A, t, p)} \frac{g_{TT}(S_A, \theta, p_r)}{g_{TT}(S_A, t, p)}.\quad (A.15.2)$$

In order to derive Eqn. (2.18.3) for α^Θ we first need an expression for $\partial\Theta/\partial T|_{S_A, p}$. This is found by differentiating with respect to *in situ* temperature the entropy equality $\eta(S_A, t, p) = \hat{\eta}(S_A, \Theta[S_A, t, p])$ obtaining

$$\frac{\partial\Theta}{\partial T} \bigg|_{S_A, p} = \eta_T(S_A, t, p) \frac{\partial\Theta}{\partial \eta} \bigg|_{S_A} = -(T_0 + \theta) \frac{g_{TT}(S_A, t, p)}{c_p^0} = \frac{(T_0 + \theta) c_p(S_A, t, p)}{(T_0 + t) c_p^0},\quad (A.15.3)$$

where the second part of this equation has used Eqn. (A.12.4) for $\Theta_\eta|_{S_A}$. This is then used to obtain the desired expression Eqn. (2.18.3) for α^Θ as follows

$$\alpha^\Theta = \frac{1}{v} \frac{\partial v}{\partial \Theta} \bigg|_{S_A, p} = \frac{1}{v} \frac{\partial v}{\partial T} \bigg|_{S_A, p} \left(\frac{\partial\Theta}{\partial T} \bigg|_{S_A, p} \right)^{-1} = - \frac{g_{TP}(S_A, t, p)}{g_P(S_A, t, p)} \frac{c_p^0}{(T_0 + \theta) g_{TT}(S_A, t, p)}.\quad (A.15.4)$$

In order to derive Eqn. (2.19.2) for β^θ we first need an expression for $\partial\theta/\partial S_A|_{T, p}$. This is found by differentiating with respect to Absolute Salinity the entropy equality $\eta(S_A, t, p) = \eta(S_A, \theta[S_A, t, p, p_r], p_r)$ which defines potential temperature, obtaining

$$\begin{aligned}
\left. \frac{\partial \theta}{\partial S_A} \right|_{T,p} &= \left. \theta_{\eta} \right|_{S_A} \left[\eta_{S_A}(S_A, t, p) - \eta_{S_A}(S_A, \theta, p_r) \right] \\
&= \frac{(T_0 + \theta)}{c_p(S_A, \theta, p_r)} \left[\mu_T(S_A, \theta, p_r) - \mu_T(S_A, t, p) \right] \\
&= \left[g_{S_A T}(S_A, t, p) - g_{S_A T}(S_A, \theta, p_r) \right] / g_{TT}(S_A, \theta, p_r),
\end{aligned} \tag{A.15.5}$$

where Eqns. (A.12.5) and (A.12.7) have been used with a general reference pressure p_r rather than with $p_r = 0$. By differentiating $\rho = \tilde{\rho}(S_A, \theta[S_A, t, p, p_r], p)$ with respect to Absolute Salinity it can be shown that (Gill (1982), McDougall (1987a))

$$\beta^{\theta} = \frac{1}{\rho} \left. \frac{\partial \rho}{\partial S_A} \right|_{\theta, p} = \frac{1}{\rho} \left. \frac{\partial \rho}{\partial S_A} \right|_{T, p} + \alpha^{\theta} \left. \frac{\partial \theta}{\partial S_A} \right|_{T, p}, \tag{A.15.6}$$

and using Eqn. (A.15.5) we arrive at the desired expression Eqn. (2.19.2) for β^{θ}

$$\beta^{\theta} = - \frac{g_{S_A P}(S_A, t, p)}{g_P(S_A, t, p)} + \frac{g_{TP}(S_A, t, p) \left[g_{S_A T}(S_A, t, p) - g_{S_A T}(S_A, \theta, p_r) \right]}{g_P(S_A, t, p) g_{TT}(S_A, t, p)}. \tag{A.15.7}$$

Note that the terms in the natural logarithm of the square root of Absolute Salinity cancel from the two parts of the square brackets in Eqns. (A.15.5) and (A.15.7).

In order to derive Eqn. (2.19.3) for β^{Θ} we first need an expression for $\partial \Theta / \partial S_A|_{T, p}$. This is found by differentiating with respect to Absolute Salinity the entropy equality $\eta(S_A, t, p) = \hat{\eta}(S_A, \Theta[S_A, t, p])$ obtaining (using Eqns. (A.12.4) and (A.12.8))

$$\begin{aligned}
\left. \frac{\partial \Theta}{\partial S_A} \right|_{T, p} &= \left. \Theta_{\eta} \right|_{S_A} \left[\eta_{S_A}(S_A, t, p) - \hat{\eta}_{S_A}|_{\Theta} \right] \\
&= \left[\mu(S_A, \theta, 0) - (T_0 + \theta) \mu_T(S_A, t, p) \right] / c_p^0 \\
&= \left[g_{S_A}(S_A, \theta, 0) - (T_0 + \theta) g_{S_A T}(S_A, t, p) \right] / c_p^0.
\end{aligned} \tag{A.15.8}$$

Differentiating $\rho = \hat{\rho}(S_A, \Theta[S_A, t, p], p)$ with respect to Absolute Salinity leads to

$$\beta^{\Theta} = \frac{1}{\rho} \left. \frac{\partial \rho}{\partial S_A} \right|_{\Theta, p} = \frac{1}{\rho} \left. \frac{\partial \rho}{\partial S_A} \right|_{T, p} + \alpha^{\Theta} \left. \frac{\partial \Theta}{\partial S_A} \right|_{T, p}, \tag{A.15.9}$$

and using Eqn. (A.15.8) we arrive at the desired expression (2.19.3) for β^{Θ} namely

$$\beta^{\Theta} = - \frac{g_{S_A P}(S_A, t, p)}{g_P(S_A, t, p)} + \frac{g_{TP}(S_A, t, p) \left[g_{S_A T}(S_A, t, p) - g_{S_A}(S_A, \theta, 0) / (T_0 + \theta) \right]}{g_P(S_A, t, p) g_{TT}(S_A, t, p)}. \tag{A.15.10}$$

Note that the terms in the natural logarithm of the square root of Absolute Salinity cancel from the two parts of the square brackets in Eqns. (A.15.8) and (A.15.10).